

task_5u1zbroaz75w39jk_with_calculation

Student Group

First Name	Surname	Matrikel Nr.

Table of Contents

Exercise E1 Electrostatics I (written test, approx. 8 % of a 120-minute written test, SS2024)	2
---	---

electrostatic, field lines, exam ee2 SS2024

Exercise E1 Electrostatics I

(written test, approx. 8 % of a 120-minute written test, SS2024)

2. What has been given to you? The charges are \$q_1 = 1 \text{ nC}\$, \$q_2 = 1 \text{ nC}\$, \$q_0 = 1 \text{ nC}\$, and \$q_4 = 1 \text{ nC}\$. The charges are placed on a grid with a spacing of \$1 \text{ mm}\$. The charges are placed at the following coordinates: \$q_1\$ at \$(1, 1)\$, \$q_2\$ at \$(2, 1)\$, \$q_0\$ at \$(3, 1)\$, and \$q_4\$ at \$(4, 0)\$. The grid is shown in the diagram below. The charges are placed on a grid with a spacing of \$1 \text{ mm}\$. The charges are placed at the following coordinates: \$q_1\$ at \$(1, 1)\$, \$q_2\$ at \$(2, 1)\$, \$q_0\$ at \$(3, 1)\$, and \$q_4\$ at \$(4, 0)\$. The grid is shown in the diagram below.

Path: \$q_0 = -1 \text{ nC}\$


- \$q_1 = -5 \text{ nC}\$

Path: \$E_4 = 5907 \text{ (from N/C)}\$

- \$\vec{F}_{01} = \left(\begin{array}{c} 917 \\ 0 \end{array} \right) \text{ (in } \mu\text{N)}\$

In the first part of the exercise, we can calculate the magnitude of the force between two charges \$q_1\$ and \$q_0\$ using Coulomb's law. The force is given by \$F = k \cdot \frac{q_1 \cdot q_0}{r^2}\$, where \$k = 8.987 \cdot 10^9 \text{ N}\cdot\text{m}^2/\text{C}^2\$. The distance between \$q_1\$ and \$q_0\$ is \$r = 2 \text{ mm} = 2 \cdot 10^{-3} \text{ m}\$. The charges are \$q_1 = 1 \text{ nC} = 1 \cdot 10^{-9} \text{ C}\$ and \$q_0 = 1 \text{ nC} = 1 \cdot 10^{-9} \text{ C}\$. The force is \$F = 8.987 \cdot 10^9 \cdot \frac{(1 \cdot 10^{-9}) \cdot (1 \cdot 10^{-9})}{(2 \cdot 10^{-3})^2} = 1.123 \cdot 10^{-4} \text{ N} = 112.3 \mu\text{N}\$. Here, the force is purely on the \$x\$-axis, so \$\vec{F}_{01} = (112.3, 0) \mu\text{N}\$.

$$|\vec{F}_{01}| = |E_4| \cdot |q_0| \rightarrow E_4 = \frac{|\vec{F}_{01}|}{|q_0|} = \frac{917 \cdot 10^{-6} \text{ N}}{1 \cdot 10^{-9} \text{ C}} = 917 \cdot 10^3 \frac{\text{V}}{\text{m}} = 917 \cdot 10^3 \frac{\text{V}}{\text{m}}$$



1. Calculate the single forces \$\vec{F}_{01}\$, \$\vec{F}_{02}\$, \$\vec{F}_{03}\$, on the charge \$q_0\$!

Path

First, calculate the magnitude of the forces, like \$\vec{F}_{01}\$.
 The force \$\vec{F}_{01}\$ is purely on the \$x\$-axis and therefore equal to \$F_{01,x}\$.

$$\vec{F}_{01} = F_{01,x} = \frac{1}{4\pi\epsilon_0} \cdot \frac{q_1 \cdot q_0}{r_{01}^2} = \frac{1}{4\pi \cdot 8.854 \cdot 10^{-12} \text{ As/Vm}} \cdot \frac{1 \cdot 10^{-9} \text{ C} \cdot 5 \cdot 10^{-9} \text{ C}}{(7 \cdot 10^{-3} \text{ m})^2} =$$

```

917.\;.\!.\!.\! \cdot 10^{-6} {\rm \{(As)^2 \cdot Vm\}\over{As \cdot m^2}} =
917.\;.\!.\!.\! \cdot 10^{-6} {\rm \{VAs\}\over{m}} = 917.\;.\!.\!.\! \cdot 10^{-6}
{\rm \{Ws\}\over{m}} \quad \&= 917.\;.\!.\!.\! {\rm \mu N} \quad \text{(to the right)}
\end{align*}

```

```

Similarly, we get for  $\vec{F}_{02}$  and  $\vec{F}_{03}$ 
\begin{align*}
\vec{F}_{02} = F_{02,x} \quad \&= -1997.\;.\!.\!.\! {\rm \mu N} \quad \text{(to the right)} \quad \& \\
\vec{F}_{03} = F_{03,y} \quad \&= -1123.\;.\!.\!.\! {\rm \mu N} \quad \text{(to the top)} \quad \& \\
\end{align*}

```

From:

<https://wiki.mexle.org/> - **MEXLE Wiki**

Permanent link:

https://wiki.mexle.org/ee2/task_5u1zbroaz75w39jk_with_calculation

Last update: **2024/07/15 15:18**

