

task_dtoqvpvrbdtozfk_with_calculation

Student Group

First Name	Surname	Matrikel Nr.

Table of Contents

Exercise E1 Electrostatics I (written test, approx. 10 % of a 120-minute written test, SS2022)	2
--	---

electrostatic, field lines, exam ee2 SS2022

Exercise E1 Electrostatics I

(written test, approx. 10 % of a 120-minute written test, SS2022)

2. What has been given to you? The charges are $q_1 = 1 \text{ nC}$, $q_2 = 2 \text{ nC}$, $q_3 = 3 \text{ nC}$, $q_4 = 4 \text{ nC}$. The value of the point charge q_0 is 1 nC . Which value needs E_4 to have to get a resulting force of 0 N on q_0 ?

Path: $q_0 = -1 \text{ nC}$

- $q_1 = -2 \text{ nC}$

Path: $E_4 = 2310.97 \text{ (n/mkV)}$

- $\vec{F}_{01} = \left(\begin{array}{c} 19.97 \\ 0 \\ 0 \end{array} \right) \text{ (nN)}$

In the x -direction the force components are $F_{01,x} = 19.97 \text{ nN}$. The force F_{02} is purely in the y -direction $F_{02,y} = 3.894 \text{ nN}$.

$$|\vec{F}_{01}| = \sqrt{F_{01,x}^2 + F_{01,y}^2} = \sqrt{19.97^2 + 0^2} = 19.97 \text{ nN}$$

$$|\vec{F}_{02}| = \sqrt{F_{02,x}^2 + F_{02,y}^2} = \sqrt{0^2 + 3.894^2} = 3.894 \text{ nN}$$

$$|\vec{F}_{03}| = \sqrt{F_{03,x}^2 + F_{03,y}^2} = \sqrt{10.05^2 + 10.05^2} = 14.20 \text{ nN}$$


$$|\vec{F}_{04}| = \sqrt{F_{04,x}^2 + F_{04,y}^2} = \sqrt{0^2 + 0^2} = 0 \text{ nN}$$

$$|\vec{F}_{01}| = |E_4| \cdot |q_0| \Rightarrow E_4 = \frac{|\vec{F}_{01}|}{|q_0|} = \frac{19.97 \text{ nN}}{1 \text{ nC}} = 19.97 \text{ nV/m}$$

$$|\vec{F}_{02}| = |E_4| \cdot |q_0| \Rightarrow E_4 = \frac{|\vec{F}_{02}|}{|q_0|} = \frac{3.894 \text{ nN}}{1 \text{ nC}} = 3.894 \text{ nV/m}$$

$$|\vec{F}_{03}| = |E_4| \cdot |q_0| \Rightarrow E_4 = \frac{|\vec{F}_{03}|}{|q_0|} = \frac{14.20 \text{ nN}}{1 \text{ nC}} = 14.20 \text{ nV/m}$$

$$|\vec{F}_{04}| = |E_4| \cdot |q_0| \Rightarrow E_4 = \frac{|\vec{F}_{04}|}{|q_0|} = \frac{0 \text{ nN}}{1 \text{ nC}} = 0 \text{ nV/m}$$



1. Calculate the single forces \vec{F}_{01} , \vec{F}_{02} , \vec{F}_{03} , on the charge q_0 !

Path

First, calculate the magnitude of the forces, like \vec{F}_{01} .

The force \vec{F}_{01} is purely on the x -axis and therefore equal to $F_{01,x}$.

$$\vec{F}_{01} = F_{01,x} \hat{x}$$

$$F_{01,x} = \frac{1}{4\pi\epsilon_0} \cdot \frac{q_1 \cdot q_0}{r_{01}^2} = \frac{1}{4\pi \cdot 8.854 \cdot 10^{-12} \text{ As/Vm}} \cdot \frac{1 \cdot 1 \cdot 10^{-9} \text{ C} \cdot 2 \cdot 10^{-9} \text{ C}}{(3 \cdot 10^{-2} \text{ m})^2} = 19.97 \dots \cdot 10^{-6} \frac{\text{As}^2 \cdot \text{Vm}}{\text{As} \cdot \text{m}^2} = 19.97 \dots$$

$$F_{02,y} = \frac{1}{4\pi\epsilon_0} \cdot \frac{q_2 \cdot q_0}{r_{02}^2} = \frac{1}{4\pi \cdot 8.854 \cdot 10^{-12} \text{ As/Vm}} \cdot \frac{2 \cdot 1 \cdot 10^{-9} \text{ C} \cdot 3 \cdot 10^{-9} \text{ C}}{(2 \cdot 10^{-2} \text{ m})^2} = 3.894 \dots \cdot 10^{-6} \frac{\text{As}^2 \cdot \text{Vm}}{\text{As} \cdot \text{m}^2} = 3.894 \dots$$

$$F_{03,x} = \frac{1}{4\pi\epsilon_0} \cdot \frac{q_3 \cdot q_0}{r_{03}^2} = \frac{1}{4\pi \cdot 8.854 \cdot 10^{-12} \text{ As/Vm}} \cdot \frac{3 \cdot 1 \cdot 10^{-9} \text{ C} \cdot 3 \cdot 10^{-9} \text{ C}}{(2 \cdot 10^{-2} \text{ m})^2} = 10.05 \dots \cdot 10^{-6} \frac{\text{As}^2 \cdot \text{Vm}}{\text{As} \cdot \text{m}^2} = 10.05 \dots$$

$$F_{03,y} = \frac{1}{4\pi\epsilon_0} \cdot \frac{q_3 \cdot q_0}{r_{03}^2} = \frac{1}{4\pi \cdot 8.854 \cdot 10^{-12} \text{ As/Vm}} \cdot \frac{3 \cdot 1 \cdot 10^{-9} \text{ C} \cdot 3 \cdot 10^{-9} \text{ C}}{(2 \cdot 10^{-2} \text{ m})^2} = 10.05 \dots \cdot 10^{-6} \frac{\text{As}^2 \cdot \text{Vm}}{\text{As} \cdot \text{m}^2} = 10.05 \dots$$

$\cdot 10^{-6} \text{ \rm \{VAs\}\over\{m\}} = 19.97... \cdot 10^{-6} \text{ \rm \{Ws\}\over\{m\}} \quad \&= 19.97... \text{ \rm \mu N} \quad \text{\text{(to the right)}} \end{align*}$

Similarly, we get for \vec{F}_{02} and \vec{F}_{03} $\begin{align*} \vec{F}_{02} = F_{02,x} \quad \&= -28.09... \text{ \rm \mu N} \quad \text{\text{(to the right)}} \quad \backslash \backslash \\ \vec{F}_{03} \quad \&= -22.47... \text{ \rm \mu N} \quad \text{\text{(to the top left)}} \quad \backslash \backslash \end{align*}$

For \vec{F}_{03} , we have to calculate the x - and y -component.

This is possible, by using the angle α between the line through q_0 and q_3 and the positive x -axis (pointing to the right).

So, α has to be between 90° and 180° . It can be calculated by:

$\begin{align*} \alpha = \arctan\left(\frac{-4\text{ cm}}{+2\text{ cm}}\right) = \pi - 1.1071... = 180^\circ - 63.4...^\circ = 116.6...^\circ \end{align*}$

Based on this, the x - and y -component is: $\begin{align*} F_{03,x} \quad \&= |\vec{F}_{03}| \cdot \cos \alpha = 10.05... \text{ \rm \mu N} \quad \text{\text{(to the left)}} \quad \backslash \backslash \\ F_{03,y} \quad \&= |\vec{F}_{03}| \cdot \sin \alpha = 20.10... \text{ \rm \mu N} \quad \text{\text{(to the top)}} \quad \backslash \backslash \end{align*}$

From:

<https://wiki.mexle.org/> - **MEXLE Wiki**

Permanent link:

https://wiki.mexle.org/ee2/task_dtoqvpvrbdcozfk_with_calculation

Last update: **2024/07/15 15:03**

