

task_dtoqvpvrbdtozfk_with_calculation

Student Group

First Name	Surname	Matrikel Nr.

Table of Contents

Exercise E2 Electrostatics I (written test, approx. 10 % of a 120-minute written test, SS2022)	2
--	---

electrostatic, field lines, exam ee2 SS2022

Exercise E2 Electrostatics I

(written test, approx. 10 % of a 120-minute written test, SS2022)

2. What has happened to the force on \$q_0\$? The picture below shows the values of the previous results. Which value needs \$E_4\$ to have to get a resulting force of \$0\$ N on \$q_0\$?

Path: $q_0 = -1 \text{ nC}$

- $q_1 = -2 \text{ nC}$

Path: $E_4 = 2310.97 \text{ (nN/m)}$

- $\vec{F}_{01} = \left(\begin{array}{c} 19.97 \\ 0 \\ 0 \end{array} \right) \text{ (nN)}$

In the previous part, we can calculate the resulting magnitude of the force on \$q_0\$ by using the superposition principle:

$$|\vec{F}_{01}| = \sqrt{\left(\sum_i F_{i,x} \right)^2 + \left(\sum_i F_{i,y} \right)^2} = \sqrt{\left(19.97 \cdot 10^{-9} \right)^2 + \left(0 \right)^2} = 19.97 \cdot 10^{-9} \text{ N}$$

Here, this force is the only one acting on \$q_0\$ from \$q_1\$ and \$q_2\$.

$$|\vec{F}_{01}| = \sqrt{\left(\sum_i F_{i,x} \right)^2 + \left(\sum_i F_{i,y} \right)^2} = \sqrt{\left(19.97 \cdot 10^{-9} \right)^2 + \left(0 \right)^2} = 19.97 \cdot 10^{-9} \text{ N}$$


$$|\vec{F}_{01}| = |E_4| \cdot |q_0| \implies E_4 = \frac{|\vec{F}_{01}|}{|q_0|} = \frac{19.97 \cdot 10^{-9} \text{ N}}{1 \cdot 10^{-9} \text{ C}} = 19.97 \cdot 10^3 \text{ V/m}$$

$$E_4 = \frac{19.97 \cdot 10^{-9} \text{ N}}{1 \cdot 10^{-9} \text{ C}} = 19.97 \cdot 10^3 \text{ V/m}$$

$$E_4 = 19.97 \cdot 10^3 \text{ V/m}$$

$$E_4 = 19.97 \cdot 10^3 \text{ V/m}$$

$$E_4 = 19.97 \cdot 10^3 \text{ V/m}$$



1. Calculate the single forces \$\vec{F}_{01}\$, \$\vec{F}_{02}\$, \$\vec{F}_{03}\$, on the charge \$q_0\$!

Path

First, calculate the magnitude of the forces, like \$\vec{F}_{01}\$.

The force \$\vec{F}_{01}\$ is purely on the \$x\$-axis and therefore equal to

$$F_{01,x} = \frac{1}{4\pi\epsilon_0} \cdot \frac{q_1 \cdot q_0}{r_{01}^2} = \frac{1}{4\pi\epsilon_0} \cdot \frac{(-2 \text{ nC}) \cdot (-1 \text{ nC})}{(1 \text{ cm})^2} = \frac{1}{4\pi\epsilon_0} \cdot \frac{2 \cdot 10^{-18} \text{ C}^2}{10^{-4} \text{ m}^2} = 19.97 \cdot 10^{-9} \text{ N}$$

$$F_{01,x} = \frac{1}{4\pi\epsilon_0} \cdot \frac{q_1 \cdot q_0}{r_{01}^2} = \frac{1}{4\pi\epsilon_0} \cdot \frac{2 \cdot 10^{-18} \text{ C}^2}{10^{-4} \text{ m}^2} = 19.97 \cdot 10^{-9} \text{ N}$$

$$F_{01,x} = \frac{1}{4\pi\epsilon_0} \cdot \frac{q_1 \cdot q_0}{r_{01}^2} = \frac{1}{4\pi\epsilon_0} \cdot \frac{2 \cdot 10^{-18} \text{ C}^2}{10^{-4} \text{ m}^2} = 19.97 \cdot 10^{-9} \text{ N}$$

$$F_{01,x} = \frac{1}{4\pi\epsilon_0} \cdot \frac{q_1 \cdot q_0}{r_{01}^2} = \frac{1}{4\pi\epsilon_0} \cdot \frac{2 \cdot 10^{-18} \text{ C}^2}{10^{-4} \text{ m}^2} = 19.97 \cdot 10^{-9} \text{ N}$$

$$F_{01,x} = \frac{1}{4\pi\epsilon_0} \cdot \frac{q_1 \cdot q_0}{r_{01}^2} = \frac{1}{4\pi\epsilon_0} \cdot \frac{2 \cdot 10^{-18} \text{ C}^2}{10^{-4} \text{ m}^2} = 19.97 \cdot 10^{-9} \text{ N}$$

$\cdot 10^{-6} \left\{ \frac{V_A}{m} \right\} = 19.97... \cdot 10^{-6} \left\{ \frac{W}{m} \right\} \quad \&= 19.97... \left\{ \mu N \right\} \quad \text{(to the right)}$

Similarly, we get for \vec{F}_{02} and \vec{F}_{03}

$\vec{F}_{02} = F_{02,x} \quad \&= -28.09... \left\{ \mu N \right\} \quad \text{(to the right)}$
 $\vec{F}_{03} \quad \&= -22.47... \left\{ \mu N \right\} \quad \text{(to the top left)}$

For \vec{F}_{03} , we have to calculate the x - and y -component.

This is possible, by using the angle α between the line through q_0 and q_3 and the positive x -axis (pointing to the right).

So, α has to be between 90° and 180° . It can be calculated by:

$\alpha = \arctan\left(\frac{-4 \text{ cm}}{+2 \text{ cm}}\right) = \pi - 1.1071... = 180^\circ - 63.4...^\circ = 116.6...^\circ$

Based on this, the x - and y -component is:

$F_{03,x} \quad \&= |\vec{F}_{03}| \cdot \cos \alpha = 10.05... \left\{ \mu N \right\} \quad \text{(to the left)}$
 $F_{03,y} \quad \&= |\vec{F}_{03}| \cdot \sin \alpha = 20.10... \left\{ \mu N \right\} \quad \text{(to the top)}$

From:

<https://wiki.mexle.org/> - **MEXLE Wiki**

Permanent link:

https://wiki.mexle.org/electrical_engineering_and_electronics/task_dtoqpvrbdtcozfk_with_calculation

Last update: **2024/07/15 15:03**

